## A Geospatial Analysis of Conflict in Afghanistan Dalton Shaver

Faculty Advisor: Professor Steve Morris

The purpose of this analysis was to determine the spatial distribution of high concentrations of conflict in Afghanistan following the 2001 invasion by NATO coalition forces. The United States government and military, as well as NATO allies parts of the country.

Figure 2 examines the spatial lag of all 328 districts. The value in each district represents the local average standardized count of conflict events in its locale. It is clear that regions with similarly high or low values are closer together, implying positive autocorrelation.

Figure 3 shows how similar each district is compared to its neighbors. Multiple clusters of similar values appear in the southern and eastern regions of the country.

Figure 4 shows nine of the total 13 clusters predicted intersect a national highway. Conflict is less likely to occur where there is a lack of major infrastructure, most notably in central Afghanistan.

240 gilinattarotsa. The contract Statescand outher NATO forces is toget answift invasion and VWLFDOOV VLJQLILFDQW . takonaves of the Jaliban, whe dispices imany in Alfabanistae in The Hynite dispitals created, funded nead the trained three Alignan what is all Aboryain committee me tated in a security force

beigween alnowarthgrantilistungente tendstatenienther Tungandine entakigise 15, 2021.

provincial capitals. This could potentially be the result of surrounding roads and

## METHODS

Highmainin the test et its utepps ala University's Georeferenced Event Dataset which 

Stable files for it es the eeg is noncentre is teleptoning by come to lieuwerd to be invensity roof Texensts Omdirated\_interestyer of casualties in a province. We can assume that the results of a global spatial autocorrelation analysis of casualties would produce similar results as seen in Histograms were used to examine the distribution of the calculated distance between this analysis. each conflict event and the nearest provincial capital, as well as the distance between each event and the nearest highway, much of which forms Afghanistan's Ring Road.

Multiple Bar Chart was used to examine the relationship between the relative frequencies of events and casualties by province.

Single Bar Chart was used to assess the distribution of events for years 2001-2019. Simple Linear Regression Modelwas used to determine the correlation between the number of casualties and conflict events in each province.

**DBSCAN Model** 

EDVHG RQ /RFDO

## CONCLUSIONS

This analysis aimed to provide insights into the spatial distribution of conflict even over 18 years of war in Afghanistan. The global and local spatial autocorrelation proved to be positive using the Morah's tatistic, showing districts with similar amounts of conflict are more likely to be clustered together. A greater number of hotspots were found in southern and eastern Afghanistan, particularly close to nat highways. Further research in analyzing the spatial distribution of casualties and t specific involvement of belligerents would provide more effective insight into where and how war was conducted in Afghanistan since 2001.